

AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions, and listings, of claims in the application:

1. (Withdrawn) A method of mediating trafficking of a target to a lysosomal compartment of a cell comprising associating with said target a molecule containing an L-T-L peptide under conditions such that said trafficking is effected, wherein said molecule is other than an FcγRIIA molecule endogenous to said cell.
2. (Withdrawn) The method according to claim 1 wherein said target is a bacterium.
3. (Cancelled).
4. (Currently Amended) ~~The method according to claim 3~~ A method of enhancing the ability of a cell to degrade a particle comprising introducing into said cell a nucleic acid sequence encoding an Fc receptor comprising an L-T-L sequence in a cytoplasmic domain thereof, said introduction being effected under conditions such that said nucleic acid sequence is expressed and said enhancement is thereby effected,
wherein said Fc receptor comprises a is FcγRIIA cytoplasmic domain ~~or a~~
modified to comprise ~~form thereof comprising~~ at least 1 additional L-T-L peptide ~~in the cytoplasmic domain thereof.~~

5. (Currently Amended) ~~The method according to claim 3~~ A method of enhancing the ability of a cell to degrade a particle comprising introducing into said cell a nucleic acid sequence encoding an Fc receptor comprising an L-T-L sequence in a cytoplasmic domain thereof, said introduction being effected under conditions such that said nucleic acid sequence is expressed and said enhancement is thereby effected,

wherein said Fc receptor comprises a γ chain cytoplasmic domain modified to comprise comprising at least one L-T-L peptide ~~in the cytoplasmic domain thereof.~~

6. (Currently Amended) The method according to claim 3 4 or 5 wherein said cell naturally expresses Fc γ RIIA.

7. (Currently Amended) The method according to claim 3 4 or 5 wherein said cell does not naturally express Fc γ RIIA.

8. (Currently Amended) The method according to claim 3 4 or 5 wherein said cell is an endothelial cell, a fibroblast, a macrophage or an epithelial cell.

9. (Currently Amended) The method according to claim 3 4 or 5 wherein said particle is a bacterium.

10. (Currently Amended) The method according to claim ~~3~~ 4 or 5 wherein said nucleic acid sequence is introduced into said cell in a liposome, a bacterium or a viral vector.

11. (Withdrawn) A method of enhancing the ability of a cell that has on its surface FcγRIIA to degrade a particle comprising contacting said cell with an agent that increases the number of FcγRIIA molecules present on the surface of said cell so that said enhancement is effected.

12. (Withdrawn) The method according to claim 11 wherein said cell is a macrophage.

13. (Withdrawn) The method according to claim 11 wherein said agent is IFN-γ or an inhibitor of IL-4.

14. (Withdrawn) A method of targeting a particle to an FcγRIIA molecule present on a cell comprising contacting said particle with a bi-specific antibody that recognizes said particle and the extracellular domain of said FcγRIIA under conditions such that said antibody binds said particle and said FcγRIIA so that said targeting is effected.

15. (Withdrawn) The method according to claim 14 wherein said particle is a microorganism.

16. (Withdrawn) A method of inducing uptake and targeting of a particle to a phagolysosome of a cell comprising contacting said particle with an IgG antibody that recognizes said particle, which antibody has associated therewith an L-T-L- containing peptide, wherein said contacting is effected under conditions such that said antibody binds said particle and said induction is effected.

17. (Withdrawn) The method according to claim 16 wherein said particle is an immune complex.

18. (Withdrawn) The method according to claim 16 wherein said particle is a microorganism.

19. (Withdrawn) A method of inducing targeting of a particle to a phagolysosome of a cell comprising contacting said particle with an L-T-L-containing peptide, wherein said contacting is effected under conditions such that said an L-T-L-containing peptide associates with said particle and said induction is thereby effected.

20. (Withdrawn) A method of inhibiting degradation of a particle by a phagolysosome of a cell comprising contacting said cell with an agent that inhibits the

function of L-T-L-containing molecule present in said cell so that said inhibition is effected.

21. (Withdrawn) The method according to claim 2 wherein said bacterium is a mycobacterium or an antibiotic resistant bacterium.

22. (Currently Amended) ~~The method according to claim 9~~ A method of enhancing the ability of a cell to degrade a particle comprising introducing into said cell a nucleic acid sequence encoding an Fc receptor comprising an L-T-L sequence under conditions such that said nucleic acid sequence is expressed and said enhancement is effected,

wherein said ~~bacterium~~ particle is a mycobacterium or an antibiotic resistant bacterium.

23. (Withdrawn) The method according to claim 15 wherein said microorganism is a mycobacterium or an antibiotic resistant bacterium.

24. (Withdrawn) The method according to claim 18 wherein said microorganism is a mycobacterium or an antibiotic resistant bacterium.

25. (Withdrawn) The method according to claim 20 wherein said particle is an IgG-coated immune complex.